

| ICD-9-CM Code   | ICD-10-CM Code(s)   |
|-----------------|---|
| <b>185</b>      | Malignant neoplasm of prostate<br><b>C61</b> Malignant neoplasm of prostate   |
| <b>600.01</b>   | Hypertrophy (benign) of prostate with urinary obstruction and other lower urinary tract symptoms (LUTS) lower urinary tract symptoms (LUTS)   |
| ICD-10-CM Codes | Two codes are required in ICD-10-CM:<br><b>N40.1</b> Enlarged prostate with lower urinary tract symptoms (LUTS); AND<br><b>N13.8</b> Other obstructive and reflux uropathy<br>** Note under N40.1 states – Use additional code for associated symptoms, when specified:<br><ul style="list-style-type: none"> <li>• Incomplete bladder emptying (<b>R39.14</b>)</li> <li>• Nocturia (<b>R35.1</b>)</li> <li>• Straining on urination (<b>R39.16</b>)</li> <li>• Urinary frequency (<b>R35.0</b>)</li> <li>• Urinary hesitancy (<b>R39.11</b>)</li> <li>• Urinary incontinence (<b>N39.4-</b>)</li> <li>• Urinary obstruction (<b>N13.8</b>)</li> <li>• Urinary retention (<b>R33.8</b>)</li> <li>• Urinary urgency (<b>R39.15</b>)</li> <li>• Weak urinary stream (<b>R39.12</b>)</li> </ul> ** Note under N13.8 states – Code first, if applicable, any causal condition, such as:<br><ul style="list-style-type: none"> <li>• Enlarged prostate (<b>N40.1</b>)</li> </ul> |
| <b>788.20</b>   | Retention of urine, unspecified   |
| ICD-10-CM Codes | <b>R33.9</b> Retention of urine, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>R33.0</b> Drug induced retention of urine<br>** Use additional code for adverse effect, if applicable, to identify drug ( <b>T36-T50</b> with fifth or sixth character 5)<br><b>R33.8</b> Other retention of urine<br>** Code first, if applicable, any causal condition, such as: enlarged prostate ( <b>N40.1</b> )  |
| <b>188.9</b>    | Malignant neoplasm of bladder, part unspecified   |
| ICD-10-CM Codes | <b>C67.9</b> Malignant neoplasm of bladder, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>C67.0</b> Malignant neoplasm of trigone of bladder<br><b>C67.1</b> Malignant neoplasm of dome of bladder<br><b>C67.2</b> Malignant neoplasm of lateral wall of bladder<br><b>C67.3</b> Malignant neoplasm of anterior wall of bladder<br><b>C67.4</b> Malignant neoplasm of posterior wall of bladder<br><b>C67.5</b> Malignant neoplasm of bladder neck<br><b>C67.6</b> Malignant neoplasm of ureteric orifice<br><b>C67.7</b> Malignant neoplasm of urachus<br><b>C67.8</b> Malignant neoplasm of overlapping sites of bladder  |
| <b>607.84</b>   | Impotence of organic origin   |
| ICD-10-CM Codes | <b>N52.9</b> Male erectile dysfunction, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>N52.01</b> Erectile dysfunction due to arterial insufficiency<br><b>N52.02</b> Corporo-venous occlusive erectile dysfunction<br><b>N52.03</b> Combined arterial insufficiency and corporo-venous occlusive erectile dysfunction   |

| ICD-9-CM Code           | ICD-10-CM Code(s)   |
|-------------------------|---|
| <b>607.84</b>           | Continued   |
| ICD-10-CM Codes (cont.) | <b>N52.1</b> Erectile dysfunction due to diseases classified elsewhere<br>** Code first underlying disease<br><b>N52.2</b> Drug-induced erectile dysfunction<br><b>N52.31</b> Erectile dysfunction following radical prostatectomy<br><b>N52.32</b> Erectile dysfunction following radical cystectomy<br><b>N52.33</b> Erectile dysfunction following urethral surgery<br><b>N52.34</b> Erectile dysfunction following simple prostatectomy<br><b>N52.39</b> Other post-surgical erectile dysfunction<br><b>N52.8</b> Other male erectile dysfunction |
| <b>592.0</b>            | Calculus of kidney  |
| ICD-10-CM Codes         | <b>N20.0</b> Calculus of kidney<br><b>N20.9</b> Urinary calculus; unspecified<br><b>N20.2</b> Calculus of kidney and ureter   |
| <b>790.93</b>           | Elevated prostate specific antigen (PSA)<br><b>R97.2</b> Elevated prostate specific antigen (PSA)   |
| <b>788.21</b>           | Incomplete bladder emptying<br><b>N20.2</b> Calculus of kidney with calculus of ureter  |
| <b>788.31</b>           | Urge incontinence<br><b>N39.41</b> Urge incontinence  |
| <b>592.1</b>            | Calculus of ureter  |
| ICD-10-CM Codes         | <b>N20.1</b> Calculus of ureter<br><b>N20.2</b> Calculus of kidney with calculus of ureter  |
| <b>596.54</b>           | Neurogenic bladder NOS  |
| ICD-10-CM Codes         | <b>N31.9</b> Neuromuscular dysfunction of bladder, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>N31.0</b> Uninhibited neuropathic bladder, not elsewhere classified<br><b>N31.1</b> Reflex neuropathic bladder, not elsewhere classified<br><b>N31.2</b> Flaccid neuropathic bladder, not elsewhere classified<br><b>N31.8</b> Other neuromuscular dysfunction of bladder<br>** Use additional code to identify any associated urinary incontinence ( <b>N39.3-N39.4-</b> )          |
| <b>189.0</b>            | Malignant neoplasm of kidney, except pelvis   |
| ICD-10-CM Codes         | <b>C64.1</b> Malignant neoplasm of right kidney, except renal pelvis<br><b>C64.2</b> Malignant neoplasm of left kidney, except renal pelvis<br><b>C64.9</b> Malignant neoplasm of unspecified kidney, except renal pelvis   |
| <b>788.41</b>           | Urinary frequency<br><b>R35.0</b> Frequency of micturition<br>** Code first, if applicable, any causal condition, such as enlarged prostate   |
| <b>599.0</b>            | Urinary tract infection, site not specified   |
| ICD-10-CM Codes         | <b>N39.0</b> Urinary tract infection, site not specified<br>** Use additional code ( <b>B95-B97</b> ) to identify infectious agent  |

| ICD-9-CM Code           | ICD-10-CM Code(s)   |
|-------------------------|---|
| <b>599.0</b>            | Continued   |
| ICD-10-CM Codes (cont.) | <p>** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N30.00</b> Acute cystitis without hematuria<br/> <b>N30.01</b> Acute cystitis with hematuria<br/> <b>N30.10</b> Interstitial cystitis (chronic) without hematuria<br/> <b>N30.11</b> Interstitial cystitis (chronic) with hematuria<br/> <b>N30.20</b> Other chronic cystitis without hematuria<br/> <b>N30.21</b> Other chronic cystitis with hematuria<br/> <b>N30.30</b> Trigonitis without hematuria<br/> <b>N30.31</b> Trigonitis with hematuria<br/> <b>N30.40</b> Irradiation cystitis without hematuria<br/> <b>N30.41</b> Irradiation cystitis with hematuria<br/> <b>N30.80</b> Other cystitis without hematuria<br/> <b>N30.81</b> Other cystitis with hematuria<br/> <b>N30.90</b> Cystitis, unspecified without hematuria<br/> <b>N30.91</b> Cystitis, unspecified with hematuria<br/> <b>N15.9</b> Renal tubule-interstitial disease, unspecified<br/> <b>N34.1</b> Nonspecific urethritis<br/> <b>N34.2</b> Other urethritis</p> |
| <b>625.6</b>            | Stress incontinence, female   |
| ICD-10-CM Codes         | <p><b>N39.3</b> Stress incontinence (female) (male)<br/>           ** Code also any associated overactive bladder (<b>N32.81</b>).</p>  |
| <b>599.71</b>           | Gross hematuria   |
| ICD-10-CM Codes         | <p><b>R31.0</b> Gross hematuria<br/> <b>R31.1</b> Benign essential microscopic hematuria<br/> <b>R31.2</b> Other microscopic hematuria<br/> <b>R31.9</b> Unspecified hematuria</p>  |
| <b>788.32</b>           | Stress incontinence, male   |
| ICD-10-CM Codes         | <p><b>N39.3</b> Stress incontinence (female) (male)<br/>           ** Code also any associated overactive bladder (<b>N32.81</b>).</p>  |
| <b>239.5</b>            | Neoplasm of unspecified nature, other genitourinary organs  |
|                         | <b>D49.5</b> Neoplasm of unspecified behavior of other genitourinary organs   |
| <b>598.9</b>            | Urethral stricture, unspecified   |
| ICD-10-CM Codes         | <p><b>N35.9</b> Urethral stricture, unspecified<br/>           ** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N35.010</b> Post-traumatic urethral stricture, male, meatal<br/> <b>N35.011</b> Post-traumatic bulbous urethral stricture<br/> <b>N35.012</b> Post-traumatic membranous urethral stricture<br/> <b>N35.013</b> Post-traumatic anterior urethral stricture<br/> <b>N35.014</b> Post-traumatic urethral stricture, male, unspecified<br/> <b>N35.021</b> Urethral stricture due to childbirth<br/> <b>N35.028</b> Other post-traumatic urethral stricture, female<br/> <b>N35.111</b> Postinfective urethral stricture, not elsewhere classified, male, meatal<br/> <b>N35.112</b> Postinfective bulbous urethral stricture, not elsewhere classified</p>   |

| ICD-9-CM Code           | ICD-10-CM Code(s)  |
|-------------------------|--|
| <b>598.9</b>            | Continued  |
| ICD-10-CM Codes (cont.) | <p><b>N35.113</b> Postinfective membranous urethral stricture, not elsewhere classified<br/> <b>N35.114</b> Postinfective anterior urethral stricture, not elsewhere classified<br/> <b>N35.119</b> Postinfective urethral stricture, not elsewhere classified, male, unspecified<br/> <b>N35.12</b> Postinfective urethral stricture, not elsewhere classified, female<br/> <b>N35.8</b> Other urethral stricture, unspecified</p>  |
| <b>605</b>              | Redundant prepuce and phimosis   |
| ICD-10-CM Codes         | <p><b>N47.0</b> Adherent prepuce, newborn<br/> <b>N47.1</b> Phimosis<br/> <b>N47.2</b> Paraphimosis<br/> <b>N47.8</b> Other disorders of prepuce</p>   |
| <b>591</b>              | Hydronephrosis   |
| ICD-10-CM Codes         | <p><b>N13.30</b> Unspecified hydronephrosis<br/>           ** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N13.1</b> Hydronephrosis with ureteral stricture, not elsewhere classified<br/> <b>N13.2</b> Hydronephrosis with renal and ureteral calculus obstruction<br/> <b>N13.39</b> Other hydronephrosis</p>   |
| <b>603.9</b>            | Hydrocele, unspecified   |
| ICD-10-CM Codes         | <p><b>N43.3</b> Hydrocele, unspecified<br/>           ** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N43.0</b> Encysted hydrocele<br/> <b>N43.1</b> Infected hydrocele<br/>           ** Use additional code (B95-B97) to identify infectious agent<br/> <b>N43.2</b> Other hydrocele</p>  |
| <b>239.4</b>            | Neoplasm of unspecified nature, bladder  |
|                         | <b>D49.4</b> Neoplasm of unspecified behavior of bladder   |
| <b>788.43</b>           | Nocturia   |
|                         | <b>R35.1</b> Nocturia  |
| <b>257.2</b>            | Other testicular hypofunction  |
|                         | <b>E29.1</b> Testicular hypofunction<br>**Use additional code for adverse effect, if applicable, to identify drug ( <b>T36-T50</b> with fifth or sixth character 5)  |
| <b>600.91</b>           | Hyperplasia of prostate, unspecified, with urinary obstruction and other lower urinary tract symptoms (LUTS)   |
| ICD-10-CM Codes         | <p>Two codes are required in ICD-10-CM:<br/> <b>N40.1</b> Enlarged prostate with lower urinary tract symptoms (LUTS); AND<br/> <b>N13.8</b> Other obstructive and reflux uropathy ***<br/>           ** Note under <b>N40.1</b> states – Use additional code for associated symptoms, when specified:</p> <ul style="list-style-type: none"> <li>• Incomplete bladder emptying (<b>R39.14</b>)</li> <li>• Nocturia (<b>R35.1</b>)</li> <li>• Straining on urination (<b>R39.16</b>)</li> </ul> |



| ICD-9-CM Code              | ICD-10-CM Code(s)  |  |
|----------------------------|--|--|
| <b>600.91</b><br>(cont.)   | Hyperplasia of prostate, unspecified, with urinary obstruction and other lower urinary tract symptoms (LUTS)   |  |
| ICD-10-CM Codes<br>(cont.) | <ul style="list-style-type: none"> <li>• Urinary frequency (<b>R35.0</b>)</li> <li>• Urinary hesitancy (<b>R39.11</b>)</li> <li>• Urinary incontinence (<b>N39.4-</b>)</li> <li>• Urinary obstruction (<b>N13.8</b>)</li> <li>• Urinary retention (<b>R33.8</b>)</li> <li>• Urinary urgency (<b>R39.15</b>)</li> <li>• Weak urinary stream (<b>R39.12</b>)</li> </ul> <p>** Note under <b>N13.8</b> states – Code first, if applicable, any causal condition, such as:</p> <ul style="list-style-type: none"> <li>• Enlarged prostate (<b>N40.1</b>)</li> </ul>  |  |
| <b>594.1</b>               | Other calculus in bladder  | <b>N21.0</b> Calculus in bladder   |
| <b>595.2</b>               | Other chronic cystitis   |  |
| ICD-10-CM Codes            | <b>N30.20</b> Other chronic cystitis without hematuria<br><b>N30.21</b> Other chronic cystitis with hematuria<br>** Use additional code to identify infectious agent ( <b>B95-B97</b> ).   |  |
| <b>601.1</b>               | Chronic prostatitis  | <b>N41.1</b> Chronic prostatitis   |
| <b>600.00</b>              | Hypertrophy (benign) of prostate without urinary obstruction and other lower urinary tract symptoms (LUTS)   | <b>N40.0</b> Enlarged prostate without lower urinary tract symptoms (LUTS) |
| <b>752.51</b>              | Undescended testis   |  |
| ICD-10-CM Codes            | <b>Q53.10</b> Unspecified undescended testicle, unilateral<br><b>Q53.11</b> Abdominal testicle, unilateral<br><b>Q53.12</b> Ectopic perineal testis, unilateral<br><b>Q53.20</b> Undescended testicle, unspecified, bilateral<br><b>Q53.21</b> Abdominal testis, bilateral<br><b>Q53.22</b> Ectopic perineal testis, bilateral<br><b>Q53.9</b> Hypospadias, unspecified  |  |
| <b>788.30</b>              | Urinary incontinence, unspecified  |  |
| ICD-10-CM Codes            | <b>R32</b> Unspecified urinary incontinence<br><b>R39.81</b> Functional urinary incontinence<br><b>F98.0</b> Non-organic origin<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>N39.3</b> Stress incontinence (female) (male)<br>** Code also associated overactive bladder ( <b>N32.81</b> )<br><b>N39.41</b> Urge incontinence<br><b>N39.42</b> Incontinence without sensory awareness<br><b>N39.43</b> Post-void dribbling<br><b>N39.44</b> Nocturnal enuresis<br><b>N39.45</b> Continuous leakage<br><b>N39.46</b> Mixed incontinence<br><b>N39.490</b> Overflow incontinence<br><b>N39.498</b> Other specified urinary incontinence<br>** Code also any associated overactive bladder ( <b>N32.81</b> ) |  |

| ICD-9-CM Code   | ICD-10-CM Code(s)   |                      |
|-----------------|---|----------------------|
| <b>595.1</b>    | Chronic interstitial cystitis   |                      |
| ICD-10-CM Codes | <b>N30.10</b> Interstitial cystitis (chronic) without hematuria **<br><b>N30.11</b> Interstitial cystitis (chronic) with hematuria **<br>** Use additional code to identify infectious agent ( <b>B95-B97</b> ).  |                      |
| <b>593.4</b>    | Other ureteric obstruction  |                      |
| ICD-10-CM Codes | <b>N13.8</b> Other obstructive and reflux uropathy<br>** Code first, if applicable, any causal condition, such as: enlarged prostate ( <b>N40.1</b> ).  |                      |
| <b>585.9</b>    | Chronic kidney disease, unspecified   |                      |
| ICD-10-CM Codes | <b>N18.9</b> Chronic kidney disease, unspecified<br>** Code first any associated:<br><ul style="list-style-type: none"> <li>• Diabetic chronic kidney disease (<b>E08.22, E09.22, E10.22, E11.22, E13.22</b>)</li> <li>• Hypertensive chronic kidney disease (<b>I12-, I13-</b>)</li> </ul> • Use additional code to identify kidney transplant status, if applicable ( <b>Z94.0</b> )<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>N18.1</b> Chronic kidney disease, stage 1<br><b>N18.2</b> Chronic kidney disease, stage 2 (mild)<br><b>N18.3</b> Chronic kidney disease, stage 3 (moderate)<br><b>N18.4</b> Chronic kidney disease, stage 4 (severe)<br><b>N18.5</b> Chronic kidney disease, stage 5<br><b>N18.6</b> End stage renal disease<br>** Use additional code to identify dialysis status ( <b>Z99.2</b> )  |                      |
| <b>788.63</b>   | Urgency of urination  |                      |
| ICD-10-CM Codes | <b>R39.15</b> Urgency of urination<br>** Code first, if applicable, any causal condition, such as enlarged prostate ( <b>N40.1</b> ).   |                      |
| <b>788.1</b>    | Dysuria   | <b>R30.0</b> Dysuria |
| <b>599.70</b>   | Hematuria, unspecified  |                      |
| ICD-10-CM Codes | <b>R31.9</b> Hematuria, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>R31.0</b> Gross hematuria<br><b>R31.1</b> Benign essential microscopic hematuria<br><b>R31.2</b> Other microscopic hematuria<br><b>N02.0</b> Recurrent and persistent hematuria with minor glomerular abnormality<br><b>N02.1</b> Recurrent and persistent hematuria with focal and segmental glomerular lesions<br><b>N02.2</b> Recurrent and persistent hematuria with diffuse membranous glomerulonephritis<br><b>N02.3</b> Recurrent and persistent hematuria with diffuse mesangial proliferative glomerulonephritis<br><b>N02.4</b> Recurrent and persistent hematuria with diffuse endocapillary proliferative glomerulonephritis<br><b>N02.5</b> Recurrent and persistent hematuria with diffuse mesangiocapillary glomerulonephritis<br><b>N02.6</b> Recurrent and persistent hematuria with dense deposit disease |                      |



| ICD-9-CM Code           | ICD-10-CM Code(s)   |
|-------------------------|---|
| <b>599.70</b>           | Continued   |
| ICD-10-CM Codes (cont.) | <b>N02.7</b> Recurrent and persistent hematuria with diffuse crescentic glomerulonephritis<br><b>N02.8</b> Recurrent and persistent hematuria with other morphologic changes<br><b>N02.9</b> Recurrent and persistent hematuria with unspecified morphologic changes  |
| <b>593.2</b>            | Cyst of kidney, acquired <b>N28.1</b> Cyst of kidney, acquired  |
| <b>599.72</b>           | Microscopic hematuria   |
| ICD-10-CM Codes         | <b>R31.1</b> Benign essential microscopic hematuria<br><b>R31.2</b> Other microscopic hematuria   |
| <b>607.1</b>            | Balanoposthitis   |
| ICD-10-CM Codes         | <b>N47.6</b> Balanoposthitis<br><b>N48.1</b> Balanitis<br>** Use additional code ( <b>B95-B97</b> ) to identify infectious agent.   |
| <b>789.00</b>           | Abdominal pain, unspecified site  |
| ICD-10-CM Codes         | <b>R10.9</b> Unspecified abdominal pain<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>R10.0</b> Acute abdomen<br><b>R10.10</b> Upper abdominal pain, unspecified<br><b>R10.11</b> Right upper quadrant pain<br><b>R10.12</b> Left upper quadrant pain<br><b>R10.13</b> Epigastric pain<br><b>R10.2</b> Pelvic and perineal pain<br><b>R10.30</b> Lower abdominal pain, unspecified<br><b>R10.31</b> Right lower quadrant pain<br><b>R10.32</b> Left lower quadrant pain<br><b>R10.33</b> Periumbilical pain<br><b>R10.84</b> Generalized abdominal pain |
| <b>752.61</b>           | Hypospadias   |
| ICD-10-CM Codes         | <b>Q54.9</b> Hypospadias, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>Q54.0</b> Hypospadias, balanic<br><b>Q54.1</b> Hypospadias, penile<br><b>Q54.2</b> Hypospadias, penoscrotal<br><b>Q54.3</b> Hypospadias, perineal<br><b>Q54.8</b> Other hypospadias   |

| ICD-9-CM Code   | ICD-10-CM Code(s)  |
|-----------------|--|
| <b>596.9</b>    | Unspecified disorder of bladder  |
| ICD-10-CM Codes | <b>N32.9</b> Bladder disorder, unspecified<br>** There are more specific code choice selections available in ICD-10-CM. These include:<br><b>N32.0</b> Bladder neck obstruction<br><b>N32.3</b> Diverticulum of bladder  |
| <b>788.36</b>   | Nocturnal enuresis <b>N39.44</b> Nocturnal enuresis  |
| <b>600.10</b>   | Nodular prostate without urinary obstruction <b>N40.0</b> Enlarged prostate without lower urinary tract symptoms (LUTS)  |
| <b>608.9</b>    | Unspecified disorder of male genital organs <b>N50.9</b> Disorder of male genital organs, unspecified  |
| <b>604.90</b>   | Orchitis and epididymitis, unspecified   |
| ICD-10-CM Codes | <b>N45.1</b> Epididymitis<br><b>N45.2</b> Orchitis<br><b>N45.3</b> Epididymo-orchitis<br>** Use additional code ( <b>B95-B97</b> ) to identify infectious agent  |
| <b>608.1</b>    | Spermatocele   |
| ICD-10-CM Codes | <b>N43.40</b> Spermatocele of epididymis, unspecified<br><b>N43.41</b> Spermatocele of epididymis, single<br><b>N43.42</b> Spermatocele of epididymis, multiple  |
| <b>607.85</b>   | Peyronie's disease <b>N48.6</b> Peyronie's disease   |
| <b>598.2</b>    | Postoperative urethral stricture   |
| ICD-10-CM Codes | <b>N99.110</b> Postprocedural urethral stricture, male, meatal<br><b>N99.111</b> Postprocedural bulbous urethral stricture<br><b>N99.112</b> Postprocedural membranous urethral stricture<br><b>N99.113</b> Postprocedural anterior urethral stricture<br><b>N99.114</b> Postprocedural urethral stricture, male, unspecified<br><b>N99.12</b> Postprocedural urethral stricture, female |